

Quantum group invariant, nonextensive quantum statistical mechanics

Marcelo R. Ubriaco*
 Laboratory of Theoretical Physics
 Department of Physics
 University of Puerto Rico
 P. O. Box 23343, Río Piedras
 PR 00931-3343, USA

We study the consequences of introducing quantum group invariance in the formalism of nonextensive quantum statistical mechanics. We find that the corresponding thermodynamical system is equivalent to a Bose-Einstein gas in the Boltzmann-Gibbs formalism with a higher critical temperature than the standard Bose-Einstein case.

I. INTRODUCTION

Since the formulation of nonextensive statistical mechanics [1] an impressive amount of related works has been published in several fields of physics, mathematics and biology [2]. This formalism is based on the generalized entropy

$$S_q = \frac{k}{q-1} \left(1 - \sum_R p_R^q \right), \quad (1)$$

where p_R is the probability for the ensemble to be in the state R and q is parameter which in principle can be any real number. The Tsallis entropy S_q possesses the properties of the Shannon's entropy except that of additivity. Given two independent systems Σ and Σ' , the entropy S_q satisfies the pseudoadditivity property

$$\frac{S_q^{\Sigma \cup \Sigma'}}{k} = \frac{S_q^{\Sigma}}{k} + \frac{S_q^{\Sigma'}}{k} + (1-q) \frac{S_q^{\Sigma}}{k} \frac{S_q^{\Sigma'}}{k}. \quad (2)$$

In addition, in several articles different authors have addressed the possibility that quantum groups could play a role in nonextensive physics [3]. Although nonextensive statistical mechanics does not embody any quantum group structure, it shares [4,5] with quantum groups the mathematical formalism of q -analysis. As shown in Ref. [4], the Tsallis entropy can be defined in terms of a q derivative acting on the probability distribution. This observation opens the possibility that nonextensive statistics could be also studied with the use of mathematical subjects as basic hypergeometric functions and series [6], and the theory of partitions [7]. However, in order to have a quantum group structure we need to include in a formalism or model a set of either operators or coordinates satisfying an algebra covariant under some kind of quantum group transformations [8,9]. This new structure introduces additional degrees of freedom, such that defining a quantum group invariant hamiltonian we can study the consequences that result from this new symmetry [10]. A recent study [11] showed that the thermodynamical behavior of a gas in the factorization approximation of nonextensive statistics differs considerably with the properties of a quantum group gas in the Boltzmann-Gibbs formulation. Therefore, it is our interest to study the simplest quantum group invariant nonextensive system. The thermodynamical system that we will study depends, in principle, on two parameters. One parameter is related to the generalization of Boltzmann-Gibbs thermodynamics to the formalism proposed by Tsallis. The quantum group parameter, relates the classical Lie group to the quantum group symmetry. Thus, loosely speaking, our density matrix is the result of a double deformation of the standard density matrix. This paper is organized as follows. In Sec. II we introduce a quantum group invariant density matrix and calculate the particle distribution for this model. With use of consistency arguments we show how the two parameters are related and their values restricted to a certain interval. In particular, we find that this system has a Bose-Einstein particle distribution with a higher critical temperature than a standard Bose-Einstein gas. In Sec. III we summarize our results.

*Electronic address: ubriaco@ltp.upr.clu.edu

II. NONEXTENSIVE STATISTICAL MECHANICS AND QUANTUM GROUP INVARIANCE

In this section we study the consequences of introducing quantum group invariance in nonextensive statistical mechanics. The density matrix $\hat{\rho}$ for nonextensive thermodynamics was reformulated in [12,13] in terms of normalized q -expectation values. The recent formalism defines the expectation values of an operator as

$$A_q = \frac{Tr \hat{A} \hat{\rho}^q}{Tr \hat{\rho}^q}, \quad (3)$$

where the density matrix in the grand canonical ensemble is given by the expression

$$\hat{\rho} = \frac{1}{Z} \left[1 - (1-q)\beta^* \left((\hat{H} - U_q) - \mu(\hat{N} - N_q) \right) \right]^{1/(1-q)}, \quad (4)$$

with the inverse temperature $\beta^* = \beta/c$ and the normalization $c = Tr \hat{\rho}^q$. The temperature $T_* = c/k\beta$ has been recently shown [14] to be the correct, physical, temperature in nonextensive thermodynamics. We wish to write the operator $\hat{H} - \mu\hat{N}$ in terms of quantum operators such that it is invariant under unitary quantum group transformations. The form of this operator can be inferred from the observation [10] that operators Φ_α and their adjoints $\bar{\Phi}_\alpha$ satisfying the quantum group algebra $SU_{q^{1/2}}(M)$ are defined by the equations

$$\Phi_\alpha = \phi_\alpha^{\dagger-1} q' \sum_{\lambda=\alpha+1}^M N_{\lambda/2} \{N_\alpha\}, \quad (5)$$

$$\bar{\Phi}_\alpha = \phi_\alpha^\dagger q' \sum_{\lambda=\alpha+1}^M N_{\lambda/2}, \quad (6)$$

where ϕ_α and $\phi_\alpha^\dagger$ are standard boson operators, N_α is the number operator and the bracket denote the q' -number $\{x\} = (1 - q'^x)/(1 - q')$. A simple algebraic manipulation shows that the simplest quantum group invariant operator $h = \sum_\alpha \bar{\Phi}_\alpha \Phi_\alpha$, can be written in terms of the q' -number operator as follows

$$h = \left\{ \sum_{\alpha=1}^M N_\alpha \right\}. \quad (7)$$

In our case, in addition to the sum over the internal (quantum group) degrees of freedom we have also a summation over the momentum. Therefore, based on the form of the operator h in Eq. (7) we propose the following operator

$$\hat{H} - \mu\hat{N} = \Lambda \left\{ \sum_{\kappa} \sum_{\alpha=1}^M \beta^* \epsilon'_{\kappa} \hat{N}_{\kappa,\alpha} \right\}, \quad (8)$$

where M is the number of internal degrees of freedom, $\epsilon'_\kappa = \epsilon_\kappa - \mu$, q' is related to the quantum group parameter and Λ is a constant to be determined. The number operator is written more explicitly as $\hat{N}_{\kappa,\alpha} = \phi_{\kappa,\alpha}^\dagger \phi_{\kappa,\alpha}$ where the operators $\phi_{\kappa,\alpha}$ and its adjoint are standard boson operators. A simple inspection shows that the operator $\hat{H} - \mu\hat{N}$ can be written as a summation over the internal degrees of freedom as follows

$$\hat{H} - \mu\hat{N} = \Lambda \sum_{\alpha=1}^M \left\{ \sum_{\kappa} \beta^* \epsilon'_{\kappa} \hat{N}_{\kappa,\alpha} \right\} q' \sum_{\lambda=\alpha+1}^M \sum_{\kappa'} \beta^* \epsilon'_{\kappa'} \hat{N}_{\kappa',\lambda}. \quad (9)$$

Since in the $q' \rightarrow 1$ limit the Eq. (9) should become the standard expression $\hat{H} - \mu\hat{N} = \sum_{\alpha} \sum_{\kappa} \epsilon'_{\kappa} \hat{N}_{\kappa,\alpha}$, we expect that the factor Λ to be a linear function of the temperature $1/\beta^*$. A representation of the quantum group operators can be obtained from the observation that Eq. (9) rewritten as a product $\sum_{\alpha} \bar{\Phi} \Phi$ define the operators Φ and the adjoints as

$$\begin{aligned} \bar{\Phi}_{\kappa,\alpha} &= \phi_{\kappa,\alpha}^\dagger q' \sum_{\lambda=\alpha+1}^M \sum_{\kappa'} (\beta^*/2) \epsilon'_{\kappa'} \hat{N}_{\kappa',\lambda} \\ \Phi_{\kappa,\alpha} &= \phi_{\kappa,\alpha}^{\dagger-1} q' \sum_{\lambda=\alpha+1}^M \sum_{\kappa'} (\beta^*/2) \epsilon'_{\kappa'} \hat{N}_{\kappa',\lambda} \left\{ \sum_{\kappa''} \beta^* \epsilon'_{\kappa''} \hat{N}_{\kappa'',\alpha} \right\}. \end{aligned} \quad (10)$$

In particular, for $M = 2$ and after redefinition of the operators Φ and $\bar{\Phi}$ as

$$\begin{aligned}\Omega_{\kappa,\alpha} &= \frac{1}{\{\beta^* \epsilon'_\kappa\}^{1/2}} \Phi_{\kappa,\alpha}, \\ \overline{\Omega}_{\kappa,\alpha} &= \frac{1}{\{\beta^* \epsilon'_\kappa\}^{1/2}} \overline{\Phi}_{\kappa,\alpha},\end{aligned}\tag{11}$$

it is simple to show that the operators Ω satisfy the $SU_{Q_\kappa}(2)$ covariant algebra

$$\begin{aligned}\Omega_{\kappa,2}\overline{\Omega}_{\kappa,2} &= 1 + Q_\kappa^2 \overline{\Omega}_{\kappa,2} \Omega_{\kappa,2} \\ \overline{\Omega}_{\kappa,1} \Omega_{\kappa,2} &= Q_\kappa^{-1} \Omega_{\kappa,2} \overline{\Omega}_{\kappa,1} \\ \Omega_{\kappa,1} \Omega_{\kappa,2} &= Q_\kappa^{-1} \Omega_{\kappa,2} \Omega_{\kappa,1} \\ \Omega_{\kappa,1} \overline{\Omega}_{\kappa,1} &= 1 + Q_\kappa^2 \overline{\Omega}_{\kappa,1} \Omega_{\kappa,1} - (1 - Q_\kappa^2) \overline{\Omega}_{\kappa,2} \Omega_{\kappa,2},\end{aligned}\tag{12}$$

where $Q_\kappa = q'^{\beta^* \epsilon'_\kappa/2}$. The equations (12) are covariant under the field redefinitions $\Omega' = T\Omega$ and $\overline{\Omega}' = \overline{\Omega} \overline{T}$ where the quantum matrix T is a 2×2 matrix given by

$$T = \begin{pmatrix} a & b \\ c & d \end{pmatrix},\tag{13}$$

with the elements $\{a, b, c, d\}$ generating the algebra

$$\begin{aligned}ab &= Q_\kappa^{-1}ba, & ac &= Q_\kappa^{-1}ca \\ bc &= cb, & dc &= Q_\kappa cd \\ db &= Q_\kappa bd, & da - ad &= (Q - Q_\kappa^{-1})bc \\ \det_{Q_\kappa} T &\equiv ad - Q_\kappa^{-1}bc = 1.\end{aligned}\tag{14}$$

The matrix $\overline{T}$ is obtained from the unitary conditions [15] $\overline{a} = d, \overline{b} = q^{-1}c$ and $q \in \mathbf{R}$. As is well known, the $SU_{Q_\kappa}(2)$ transformation T matrix and the corresponding R matrix

$$R = \begin{pmatrix} Q_\kappa & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & Q_\kappa - Q_\kappa^{-1} & 1 & 0 \\ 0 & 0 & 0 & Q_\kappa \end{pmatrix}$$

satisfy the algebraic relations [16]

$$RT_1T_2 = T_2T_1R, \tag{15}$$

and

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}, \tag{16}$$

where $T_1 = T \otimes 1$ and $T_2 = 1 \otimes T \in V \otimes V$, $(R_{23})_{ijk,i'j'k'} = \delta_{ii'} R_{jk,j'k'} \in V \otimes V \otimes V$. We should remark that quantum group covariance of the operator algebra is satisfied for those operators Ω with the same momentum κ . For example, for $\kappa \neq \kappa'$ we have the relation

$$\overline{\Omega}_{\kappa',1} \overline{\Omega}_{\kappa,2} = Q_\kappa \overline{\Omega}_{\kappa,2} \overline{\Omega}_{\kappa',1}, \tag{17}$$

which is not quantum group covariant [17]. For $M > 2$, and hereafter omitting the κ index, the Eq. (12) are generalized to the set of equations

$$\Omega_\alpha \overline{\Omega}_\lambda = \delta_{\alpha\lambda} + Q R_{\eta\lambda\alpha\gamma} \overline{\Omega}_\gamma \Omega_\eta \tag{18}$$

$$\Omega_\alpha \Omega_\lambda = Q^{-1} R_{\gamma\eta\lambda\alpha} \Omega_\gamma \Omega_\eta, \tag{19}$$

The $M^2 \times M^2$ matrix $R_{\alpha\lambda\gamma\eta}$ is well known in the quantum group literature [8], and in terms of our parameter Q_κ is explicitly written

$$R_{\alpha\lambda\gamma\eta} = \delta_{\alpha\gamma}\delta_{\lambda\eta}(1 + (Q - 1)\delta_{\alpha\lambda}) + (Q - Q^{-1})\delta_{\lambda\gamma}\delta_{\alpha\eta}\theta(\alpha - \lambda), \quad (20)$$

where $\theta(\alpha - \lambda) = 1$ for $\alpha > \lambda$ and zero otherwise. From Eqs. (9) and (10) we see that the quantum group invariant density matrix can be simply written as

$$\hat{\rho} = \frac{1}{Z} \left[1 - (1 - q)\beta^* \left(\sum_{\alpha=1}^M \Lambda \Phi_{\alpha} \Phi_{\alpha} + \mu N_q - U_q \right) \right]^{1/(1-q)}. \quad (21)$$

Eq. (21) is the simplest quantum group invariant density matrix one can write. It represents, after all, the density matrix for a set of free quantum group bosons Φ . By choosing the constant Λ to have the value

$$\Lambda = \frac{1 - q'}{(1 - q)\beta^*} - (1 - q')(\mu N_q - U_q), \quad (22)$$

we find that the density matrix satisfies a simple relation with boson operators

$$\hat{\rho}^q \phi_{\kappa,\alpha}^{\dagger} = \phi_{\kappa,\alpha}^{\dagger} q'^{q\beta^*} \epsilon'_{\kappa}/(1-q) \hat{\rho}^q. \quad (23)$$

Since other than integer powers of the operators ϕ and $\phi^{\dagger}$ are meaningless, the operator $\hat{\rho}^q$ only makes sense for those values of the nonextensive parameter q such that $q/(1-q)$ is a positive integer. Therefore, the parameter is restricted to the values $q = n/n+1$, for $n = 1, 2, 3, \dots$. The density matrix in Eq. (23) depends on two parameters. The parameter Q_{κ} labels, for a given value of κ , the deformation from a classical $SU(M)$ to a quantum $SU_{Q_{\kappa}}(M)$ symmetry. The parameter q indicates the departure from the Boltzmann-Gibbs formalism to nonextensive thermodynamics. A simple inspection of the operator $\hat{H} - \mu \hat{N}$ for $q' \rightarrow 1$ shows that it has the correct limit $\sum_{\alpha} \sum_{\kappa} \epsilon'_{\kappa} N_{\kappa,\alpha}$ if the two parameters are related according to

$$1 - q' = (1 - q)f(q), \quad (24)$$

where $f(q) \rightarrow 1$ as $q \rightarrow 1$. This relation implies that we cannot take the two limits $q' \rightarrow 1$ and $q \rightarrow 1$ independently without avoiding the trivial solution $f(q) = 0$. In addition, according to Eq. (24) the quantum group parameter q' is also restricted to the interval $0 < q' \leq 1$. We consider for simplicity and without loss of generality the case $q = q'$. The average number of particles with energy ϵ_{κ} is given by

$$\begin{aligned} n_{\kappa} &= Tr \hat{\rho}^q \phi_{\kappa}^{\dagger} \phi_{\kappa} / Tr \hat{\rho}^q \\ &= \frac{M}{q^{-q\beta^*} \epsilon'_{\kappa}/(1-q) - 1}, \end{aligned} \quad (25)$$

which is functionally equivalent to the Bose-Einstein distribution in Boltzmann-Gibbs thermodynamics. It is simple to check that the correct limit of the Bose-Einstein distribution $n_{\kappa} = 1/(exp(\beta\epsilon'_{\kappa}) - 1)$ is obtained for $q = 1$. Therefore, the particle distribution is equivalent to the Bose-Einstein textbook case

$$n_{\kappa} = \frac{M}{e^{\beta'^* (\epsilon_{\kappa} - \mu)} - 1}, \quad (26)$$

where $\beta'^* = -\beta^* q \ln q/(1-q)$ and the fugacity $z = exp(\beta'^* \mu)$. With use of some algebra we find that the normalization constant $c = Tr \hat{\rho}^q$ is given by the equation

$$c = \frac{1}{Z^q} Tr \left[q \sum_{\kappa} \sum_{\alpha=1}^M \beta^* \epsilon'_{\kappa} \hat{N}_{\kappa,\alpha} (1 - (1 - q)\beta^* (\mu N_q - U_q)) \right]^{q/(1-q)}, \quad (27)$$

leading to

$$c = \frac{1}{Z^q} [1 - (1 - q)\beta^* (\mu N_q - U_q)]^{q/(1-q)} \prod_{\kappa} \frac{1}{1 - q^{q\beta^*} M \epsilon'_{\kappa}/(1-q)}. \quad (28)$$

Taking into account that the constant $c = Z^{1-q}$ [13] we obtain that

$$\ln c = q \ln [1 - (1 - q)\beta^* (\mu N_q - U_q)] - (1 - q) \sum_{\kappa} \ln \left(1 - q^{q\beta^*} M \epsilon'_{\kappa}/(1-q) \right). \quad (29)$$

The summation over κ can be approximated, as usual, by an integral with the result

$$c = [1 - (1 - q)\beta^*(\mu N_q - U_q)]^q \exp \left[\frac{-4V(1 - q)^{5/2}}{(qM|\ln q|)^{3/2}\sqrt{\pi}\lambda'^{*3}} g_{5/2}(z^M) \right], \quad (30)$$

where the Bose-Einstein function $g_{5/2}(x) = \sum_{n=1}^{\infty} x^n/n^{5/2}$ and the thermal wavelength $\lambda^* = h\sqrt{\beta^*/2\pi m}$. Since the particle distribution in Eq. (26) is of the Bose-Einstein type this system will exhibit Bose-Einstein condensation but with a different critical temperature than the textbook case.

The results for the total average number of particles N_q and the total energy $U_q = Tr \sum_{\kappa} \sum_{\alpha} \epsilon_{\kappa} \phi_{\kappa}^{\dagger} \phi_{\kappa} \hat{\rho}^q / c$, can be readily copied from the standard results with the replacement of β by β'^*

$$N_q = N_0 + \frac{MV}{\lambda'^{*3}} g_{3/2}(z), \quad (31)$$

$$U_q = \frac{3}{2} \frac{MV}{\beta'^* \lambda'^{*3}} g_{5/2}(z), \quad (32)$$

with $\lambda'^* = h\sqrt{\beta'^*/2\pi m}$. The critical temperature is therefore given by the expression

$$T_c^* = \frac{-(1 - q)}{q \ln q} T_c^{BE}, \quad (33)$$

where T_c^{BE} is the critical temperature for a Bose-Einstein gas. Since the factor in Eq. (33) is $\frac{-(1-q)}{q \ln q} > 1$ we find that for the interval $0 < q < 1$ the critical temperature satisfies that $T_c^* > T_c^{BE}$. Figure 1 is a graph that shows the dependence of the critical temperature T_c^* on the parameter q .




FIG. 1. The dependence of T_c^*/T_c^{BE} on the parameter q where T_c^* is the critical temperature of our thermodynamic system and T_c^{BE} is the critical temperature for $q = 1$. The dots indicate the allowed values of q .

III. CONCLUSIONS

In this paper we have been able to define a quantum group invariant density matrix in nonextensive quantum statistical mechanics. For a given momentum κ the quantum group operators satisfy algebraic relations covariant under $SU_{q'\beta^*\epsilon'_{\kappa}/2}(M)$ transformations. Although the two deformations in the density matrix are originally unrelated, we found that the parameters q' and q are related through the equation $q' = 1 - (1 - q)f(q)$, where $f(q)$ is an arbitrary function satisfying $f(q = 1) = 1$ and the nonextensive parameter q takes values $q = n/n + 1, n = 1, 2, 3, \dots$. A simple calculation showed that the particle distribution function is equivalent to the standard Bose-Einstein distribution but

the critical temperature is higher than the critical temperature of the standard case. Thus, according to the results reported in this paper, the thermodynamics of a quantum group invariant system in nonextensive thermodynamics is equivalent to the thermodynamical behavior of a system of free bosons in the extensive, Boltzmann-Gibbs, formulation.

- [1] C. Tsallis, J. Stat. Phys. **52**, 479 (1988). E. Curado and C. Tsallis, J. Phys. A **24**, L69 (1991); **24**, 3187(E) (1991); **25**, 1019(E) (1992).
- [2] A list of works can be found in tsallis.cat.cbpf.br/biblio.htm.
- [3] C. Tsallis, Phys. Lett. A **195**, 329 (1994); S. Abe, *ibid.* **224**, 326 (1997); *ibid.* **244**, 229 (1998).
- [4] S. Abe, Phys. Lett. A **224**, 326 (1997).
- [5] R. Johal, Phys. Rev. E **58**, 4147 (1998).
- [6] See, for example, H. Exton, *q-Hypergeometric Functions and Applications*, (E. Horwood Ltd., Chichester, 1983); N. J. Fine *Basic Hypergeometric Series and Applications* (Am. Math. Soc., Providence, 1988).
- [7] G. E. Andrews, *Theory of Partitions*, (Addison-Wesley, Reading, 1976).
- [8] J. Wess and B. Zumino, Nucl. Phys. B (Proc. Suppl.) **18**, 302 (1990); M. Jimbo ed., *Yang- Baxter equation in integrable systems*, Advanced series in Mathematical Physics V.10 (World Scientific, 1990); V. Chari and A. Pressley, *A guide to Quantum Groups*, (Cambridge Univ. Press, 1994); J. C. Jantzen, *Lectures on Quantum Groups*, Graduate Studies in Mathematics V. 6 (American Mathematical Society, 1996); S. Majid, *Foundations of Quantum Group Theory*, (Cambridge Univ. Press, 1996), and references therein.
- [9] S. L. Woronowicz, Publ. Res. Ins. Math. Sci. **23**, 117 (1987); Yu Manin, Commun. Math. Phys. **123**, 163 (1989).
- [10] M. R. Ubriaco, Phys. Rev. E **55**, 291 (1997); *ibid* **57**, 179 (1998).
- [11] M. R. Ubriaco, Phys. Rev. E **60**, 165 (1999).
- [12] C. Tsallis, R. S. Mendes and A.R. Plastino, Physica A **261**, 534 (1998).
- [13] E. K. Lenzi, R. S. Mendes and A.K. Rajagopal, Quantum Statistical Mechanics for Nonextensive Systems II, cond-mat/9904100.
- [14] S. Abe, S. Martínez, F. Pennini and A. Plastino, Nonextensive thermodynamic relations, cond-mat/0011012.
- [15] S. Vokos, J. Wess and B. Zumino, *Symmetry in Nature* (Scuola Normale Superiore Publ., Pisa, Italy, 1989).
- [16] L. A. Takhtajan, Adv. Stud. Pure Math. **19**, 1 (1989), and references therein.
- [17] M. R. Ubriaco, Mod. Phys. Lett. A **8**, 2213 (1993); **10**, 2223(E) (1995).